

Exclusive production of $f_1(1285)$ meson at low and high energies

Piotr Lebiedowicz (IFJ PAN)

in collaboration with

Antoni Szczurek (IFJ PAN)

Piotr Salabura (Jagiellonian U.)

Otto Nachtmann (Heidelberg U.)

Josef Leutgeb (TU Wien)

Anton Rebhan (TU Wien)



THE HENRYK NIEWODNICZAŃSKI
INSTITUTE OF NUCLEAR PHYSICS
POLISH ACADEMY OF SCIENCES

Contents

1. Introduction
2. Formalism
3. Selected results
4. Conclusions

MESON 2021

16th International Workshop on Meson Physics

17th - 20th May 2021

Introduction | Motivation

In this talk we will be concerned with **central exclusive production (CEP)** of axial-vector $f_1(1285)$ meson in proton-(anti)proton collisions at c.m. energies:

- low: HADES (pp) and PANDA ($p\bar{p}$) at FAIR ← *Lebiedowicz, Nachtmann, Salapura, Szczurek, arXiv:2105.07192*
- intermediate: WA102, COMPASS
- high: RHIC, LHC ← *Lebiedowicz, Leutgeb, Nachtmann, Rebhan, Szczurek, PRD 102 (2020) 114003*

The $f_1(1285)$ meson was measured

- in two-photon interactions in e^+e^- reactions (MARKII, TPC/Two-Gamma, L3)
see: A. Szczurek, PRD 102 (2020) 113015 ← on production of f_1 mesons at e^+e^- collisions with double-tagging as a way to constrain the axial meson light-by-light contribution to the muon g-2 and hyperfine splitting of muonic hydrogen
- in the photoproduction process $\gamma p \rightarrow f_1 p$ (CLAS Collaboration)
- in proton-proton collisions for c.m. energies 12.7 and 29.1 GeV (WA102) and for 13 TeV at the LHC (ATLAS)
[R. Sikora, CERN-THESIS-2020-235]


Why study the $pp \rightarrow pp f_1$ process?

- **What is underlying production mechanism for studies of f_1 CEP** at near threshold and at LHC?
 - Poorly known VVf_1 and $IPIPf_1$ coupling strengths and the corresponding vertex form factors
 - Can it be described in holographic QCD?
- **What is underlying decay mechanism?**
 - e.g., $f_1(1285) \rightarrow 4\pi$ decay via $\rho\rho$ or/and $\pi a_1(1260) (\rightarrow \rho\pi)$
 - transition form factors e.g., $\gamma^*\gamma^* \rightarrow f_1$, $f_1 \rightarrow \gamma\gamma^* \rightarrow \gamma e^+e^-$
 - What is the nature of the $f_1(1285)$? For instance, is it a normal $q\bar{q}$ state or $\bar{K}K^*$ molecule?
see: Aceti, Dias, Oset, EPJA 51 (2015) 48; Aceti, Xie, Oset, PLB 750 (2015) 609
- **What is optimal observation channel of the $f_1(1285)$?**

VV-fusion mechanism

$$p(p_a, \lambda_a) + p(p_b, \lambda_b) \rightarrow p(p_1, \lambda_1) + f_1(k, \lambda_{f_1}) + p(p_2, \lambda_2)$$

$p_{a,b}, p_{1,2}$ and $\lambda_{a,b}, \lambda_{1,2} = \pm\frac{1}{2}$: the four-momenta and helicities of protons
 k and $\lambda_{f_1} = 0, \pm 1$: the four-momentum and helicity of the f_1 meson



$$q_1 = p_a - p_1, \quad q_2 = p_b - p_2, \quad k = q_1 + q_2$$

$$t_1 = q_1^2, \quad t_2 = q_2^2, \quad m_{f_1}^2 = k^2$$

$$s = (p_a + p_b)^2 = (p_1 + p_2 + k)^2, \text{ c.m. energy squared}$$

$$s_1 = (p_1 + k)^2, \quad s_2 = (p_2 + k)^2$$

VV-fusion amplitude: $\mathcal{M}_{pp \rightarrow pp f_1}^{(VV \text{ fusion})} = \mathcal{M}_{pp \rightarrow pp f_1}^{(\rho\rho \text{ fusion})} + \mathcal{M}_{pp \rightarrow pp f_1}^{(\omega\omega \text{ fusion})}$

$$\begin{aligned} \mathcal{M}_{\lambda_a \lambda_b \rightarrow \lambda_1 \lambda_2 \lambda_{f_1}}^{(VV \text{ fusion})} &= (-i) (\epsilon^\alpha(\lambda_{f_1}))^* \bar{u}(p_1, \lambda_1) i\Gamma_{\mu_1}^{(Vpp)}(p_1, p_a) u(p_a, \lambda_a) \\ &\times i\tilde{\Delta}^{(V)\mu_1\nu_1}(s_1, t_1) i\Gamma_{\nu_1\nu_2\alpha}^{(VVf_1)}(q_1, q_2) i\tilde{\Delta}^{(V)\nu_2\mu_2}(s_2, t_2) \\ &\times \bar{u}(p_2, \lambda_2) i\Gamma_{\mu_2}^{(Vpp)}(p_2, p_b) u(p_b, \lambda_b) \end{aligned}$$

$$i\Gamma_{\mu}^{(Vpp)}(p', p) = -i\Gamma_{\mu}^{(V\bar{p}\bar{p})}(p', p) = -ig_{Vpp} F_{VNN}(t) \left[\gamma_{\mu} - i \frac{\kappa_V}{2m_p} \sigma_{\mu\nu} (p - p')^{\nu} \right]$$

$$g_{ppp} = 3.0, \quad \kappa_{\rho} = 6.1, \quad g_{\omega pp} = 9.0, \quad \kappa_{\omega} = 0$$

κ_V : tensor-to-vector coupling ratio, $\kappa_V = f_{VNN}/g_{VNN}$

$$F_{VNN}(t) = \frac{\Lambda_{VNN}^2 - m_V^2}{\Lambda_{VNN}^2 - t}$$

For the proton-antiproton collisions we have

$$\begin{aligned} \bar{u}(p_2, \lambda_2) i\Gamma_{\mu_2}^{(Vpp)}(p_2, p_b) u(p_b, \lambda_b) &\rightarrow \bar{v}(p_b, \lambda_b) i\Gamma_{\mu_2}^{(V\bar{p}\bar{p})}(p_2, p_b) v(p_2, \lambda_2) \\ &= -\bar{u}(p_2, \lambda_2) i\Gamma_{\mu_2}^{(Vpp)}(p_2, p_b) u(p_b, \lambda_b) \end{aligned}$$

$$\mathcal{M}_{p\bar{p} \rightarrow p\bar{p} M}^{(VV \text{ fusion})} = -\mathcal{M}_{pp \rightarrow pp M}^{(VV \text{ fusion})}$$

The standard form of the vector-meson propagator:

$$i\Delta_{\mu\nu}^{(V)}(k) = i \left(-g_{\mu\nu} + \frac{k_{\mu}k_{\nu}}{k^2 + i\epsilon} \right) \Delta_T^{(V)}(k^2) - i \frac{k_{\mu}k_{\nu}}{k^2 + i\epsilon} \Delta_L^{(V)}(k^2)$$

$$\Delta_T^{(V)}(t) = (t - m_V^2)^{-1}$$

For higher values of s_1 and s_2 we must take into account **reggeization**:

$$\Delta_T^{(V)}(t_i) \rightarrow \tilde{\Delta}_T^{(V)}(s_i, t_i) = \Delta_T^{(V)}(t_i) \left(\exp(i\phi(s_i)) \frac{s_i}{s_{\text{thr}}} \right)^{\alpha_V(t_i) - 1}$$

$$\phi(s_i) = \frac{\pi}{2} \exp\left(\frac{s_{\text{thr}} - s_i}{s_{\text{thr}}} \right) - \frac{\pi}{2}$$

where s_{thr} is the lowest value of s_i possible here: $s_{\text{thr}} = (m_p + m_{f_1})^2$

We use the linear form for the vector meson Regge trajectories :

$$\alpha_V(t) = \alpha_V(0) + \alpha'_V t, \quad \alpha_V(0) = 0.5, \quad \alpha'_V = 0.9 \text{ GeV}^{-2}$$

VV f_1 coupling

$$\mathcal{L}'_{VVf_1}(x) = \frac{1}{M_0^4} g_{VVf_1} (V_{\kappa\lambda}(x) \overleftrightarrow{\partial}_{\mu} \overleftrightarrow{\partial}_{\nu} V_{\rho\sigma}(x)) (\partial_{\alpha} U_{\beta}(x) - \partial_{\beta} U_{\alpha}(x)) g^{\kappa\rho} g^{\mu\sigma} \varepsilon^{\lambda\nu\alpha\beta}$$

where $V_{\kappa\lambda}(x) = \partial_{\kappa} V_{\lambda}(x) - \partial_{\lambda} V_{\kappa}(x)$, $U_{\alpha}(x)$ and $V_{\kappa}(x)$ are the fields of the f_1 meson and the vector meson V , respectively.

$M_0 \equiv 1 \text{ GeV}$ and g_{VVf_1} a dimensionless coupling constant.





$$\begin{aligned} i\Gamma_{\mu\nu\alpha}^{(VVf_1)}(q_1, q_2) &= \frac{2g_{VVf_1}}{M_0^4} [(q_1 - q_2)^{\rho} (q_1 - q_2)^{\sigma} \varepsilon_{\lambda\sigma\alpha\beta} k^{\beta} \\ &\times (q_{1\kappa} \delta_{\mu}^{\lambda} - q_1^{\lambda} g_{\kappa\mu}) (q_2^{\kappa} g_{\rho\nu} - q_{2\rho} \delta_{\nu}^{\kappa}) + (q_1 \leftrightarrow q_2, \mu \leftrightarrow \nu)] \\ &\times F^{(VVf_1)}(q_1^2, q_2^2, k^2) \end{aligned}$$

satisfies gauge invariance relations: $\Gamma_{\mu\nu\alpha}^{(VVf_1)}(q_1, q_2) q_1^{\mu} = 0, \Gamma_{\mu\nu\alpha}^{(VVf_1)}(q_1, q_2) q_2^{\nu} = 0$

$$F^{(VVf_1)}(q_1^2, q_2^2, m_{f_1}^2) = \tilde{F}_V(q_1^2) \tilde{F}_V(q_2^2) F(m_{f_1}^2) = \frac{\Lambda_V^4}{\Lambda_V^4 + (t_1 - m_V^2)^2} \frac{\Lambda_V^4}{\Lambda_V^4 + (t_2 - m_V^2)^2}$$

with $F(m_{f_1}^2) = 1$

Results



- The $\rho\rho f_1$ coupling constant is extracted from the radiative decay rate $f_1 \rightarrow \rho^0\gamma$ using the VMD approach.


from PDG : $\Gamma(f_1(1285) \rightarrow \gamma\rho^0) = 1384.7^{+305.1}_{-283.1}$ keV

from CLAS : $\Gamma(f_1(1285) \rightarrow \gamma\rho^0) = (453 \pm 177)$ keV ← we use

We consider decay $f_1 \rightarrow \rho^0\gamma \rightarrow \pi^+\pi^-\gamma$ taking ρ^0 mass distribution. We estimate the cutoff parameter Λ_ρ in the $f_1\rho\rho$ form factor:

$$F_{\rho\rho f_1}(k_\rho^2, k_\gamma^2, k^2) = F_{\rho\rho f_1}(k_\rho^2, 0, m_{f_1}^2) = \tilde{F}_\rho(k_\rho^2)\tilde{F}_\rho(0)F(m_{f_1}^2) = \tilde{F}_\rho(k_\rho^2)\tilde{F}_\rho(0)$$

- Photoproduction process:



- We assume $g_{\omega\omega f_1} = g_{\rho\rho f_1}$ based on arguments from the quark model and VMD. We assume $\Lambda_\rho = \Lambda_\omega = \Lambda_V$ and $\Lambda_{\rho NN} = \Lambda_{\omega NN} = \Lambda_{VNN}$.
- Reggeization effect included
- The t-channel V-exchange mechanism play a crucial role in reproducing the forward-peaked angular distributions, especially at higher energies. From the comparison of differential cross sections to the CLAS data we estimate:

(C7) : $\Lambda_{VNN} = 1.35$ GeV for $\Lambda_V = 0.65$ GeV, $|g_{VVf_1}| = 20.03$

(C9) : $\Lambda_{VNN} = 1.01$ GeV for $\Lambda_V = 0.8$ GeV, $|g_{VVf_1}| = 12.0$

(C10) : $\Lambda_{VNN} = 0.9$ GeV for $\Lambda_V = 1.0$ GeV, $|g_{VVf_1}| = 8.49$

(C11) : $\Lambda_{VNN} = 0.834$ GeV for $\Lambda_V = 1.5$ GeV, $|g_{VVf_1}| = 6.59$



(C11) is excluded due to small Λ_{VNN} , we stay with (C7) - (C10)

- Missing N^* resonances and s/u-channel proton exchange
- Possible $N(2300)$ contribution

→ postulated in *Y.-Y. Wang et al., PRD 95 (2017) 096015*

CLAS data:
R. Dickson et al. (CLAS Collaboration), PRC 93 (2016) 065202

Results



- No data for the $pp \rightarrow pp f_1$ and $p\bar{p} \rightarrow p\bar{p} f_1$ reactions
- High sensitivity of the VV-fusion cross section to the different sets of parameters.

In our procedure of extracting the coupling constants and the form-factor cutoff parameters from the CLAS data the dominant sensitivity is on coupling constants not on the form factors

- Reggeization effect included
- The NN FSI effect can be neglected
- We predict for the VV-fusion mechanism:







HADES: $\sqrt{s} = 3.46$ GeV
 PANDA: $\sqrt{s} = 2.25 - 5.47$ GeV

$\sigma(pp \rightarrow pp f_1) = 0.04 - 0.15 \mu\text{b}$ for $\sqrt{s} = 3.46$ GeV
 $\sigma(p\bar{p} \rightarrow p\bar{p} f_1) = 0.41 - 2.07 \mu\text{b}$ for $\sqrt{s} = 5.0$ GeV **10 x larger**

- Diffractive contribution (IPIP fusion) is very small for the HADES and PANDA energy range
 → IPIP-fusion contribution should be considered as upper limit of the cross section.
 If at the WA102 c.m. energy (29.1 GeV) there are important contributions from subleading reggeon exchanges, the IPIP contribution could be smaller (by a factor of up to 4)


Results

$\sqrt{s} = 3.46$ GeV (top) and 5.0 GeV (bottom)






θ_M is the angle between \vec{k} and \vec{p}_a

- At near threshold energy (HADES) the values of small $|t_1|$ and $|t_2|$ are not accessible kinematically
- HADES and PANDA experiments have a good opportunity to study physics of large four-momentum transfer squared \rightarrow probes corresponding form factors at relatively large values of $|t_{1,2}|$
- $\rho^0\rho^0$ - and $\omega\omega$ -fusion processes have different kinematic dependences. Both terms play similar role. But with increasing c.m. energy the averages of $|t_{1,2}|$ decrease (damping by form factors), hence the $\omega\omega$ term becomes more important
- We predict a strong preference for the outgoing nucleons to be produced with their transverse momenta being back-to-back, $d\sigma/d\phi_{pp}$ at $\phi_{pp} = \pi$



Results

Optimal observation channel of $f_1(1285)$



Simulations for HADES experiment for $\sqrt{s} = 3.46 \text{ GeV}$ using PLUTO MC generator:

$$pp \rightarrow pp\pi^+\pi^-\pi^+\pi^-$$

$$\mathcal{BR}(f_1(1285) \rightarrow \pi^+\pi^-\pi^+\pi^-) = (10.9 \pm 0.6) \%$$

$$\sigma_{back}^{4\pi} \sim 227 \mu\text{b} [1], \quad \sigma_{f_1}^{4\pi} = 16 \text{ nb}$$

Difficult to see f_1 peak on the 4π background without additional cuts

$$pp \rightarrow pp\pi^+\pi^-\pi^+\pi^-\pi^0$$

$$\mathcal{BR}(f_1(1285) \rightarrow \pi^+\pi^-\eta) = (35 \pm 15) \%$$

$$\mathcal{BR}(\eta \rightarrow \pi^+\pi^-\pi^0) = (22.92 \pm 0.28) \%$$

The narrow width of the η meson allows to set a mass cut on the $\pi^+\pi^-\pi^0$ invariant mass and suppresses the multi-pion background efficiently

Contributions and cross sections used in the simulations of the reaction $pp \rightarrow pp\pi^+\pi^-\pi^+\pi^-\pi^0$

Contribution	Cross section (μb)	
1	88	$\sigma = (88 \pm 14) \mu\text{b} [1], P = 5.5 \text{ GeV}/c$
2	0.18	estimates via double N^* production (via π^0 exchange) $pp \rightarrow N(1440)N(1535)$ and $pp \rightarrow N(1535)N(1535)$
3	0.07	$\sigma = (0.09 \pm 0.03) \mu\text{b} [2]$ for $pp \rightarrow pp\pi^+\pi^-\omega$ at $P = 6.92 \text{ GeV}/c$
4	0.012	$\sigma = (3.2 - 12.4) \text{ nb}$, see (C7) –(C10), from the $VV \rightarrow f_1$ fusion mechanism

[1] G. Alexander et al., Phys. Rev. 154 (1967) 1284

[2] S. Danieli et al., Nucl. Phys. B27 (1971) 157


The study of $f_1(1285)$ production at HADES should be feasible !

Pomeron-Pomeron fusion mechanism

At high energies double pomeron (IP) exchange is dominant production mechanism of the $f_1(1285)$

see: *Lebiedowicz, Leutgeb, Nachtmann, Rebhan, Szczurek, PRD 102 (2020) 114003*

$$p(p_a) + p(p_b) \rightarrow p(p_1) + f_1(k) + p(p_2)$$



We treat our reaction in the [tensor-pomeron approach](#)

[*Ewerz, Maniatis, Nachtmann, Ann. Phys. 342 (2014) 31*]

The pomeron and the charge conjugation $C=+1$ reggeons are described as effective rank 2 symmetric tensor exchanges. The odderon and the $C=-1$ reggeons are described as effective vector exchanges.

This approach has a good basis from nonperturbative QCD considerations.

The IP exchange can be understood as a coherent sum of exchanges of spin $2+4+6+ \dots$

[*Nachtmann, Ann. Phys. 209 (1991) 436*]

A tensor character of the pomeron is also preferred in holographic QCD, see e.g.,


Brower, Polchinski, Strassler, Tan, JHEP 12 (2007) 005

Domokos, Harvey, Mann, PRD 80 (2009) 126015

Iatrakis, Ramamurti, Shuryak, PRD 94 (2016) 045005

The Born-level amplitude within the tensor-pomeron approach:

$$\begin{aligned} \mathcal{M}_{\lambda_a \lambda_b \rightarrow \lambda_1 \lambda_2 \lambda_{f_1}}^{\text{Born}} &= (-i) (\epsilon^\mu(\lambda_{f_1}))^* \bar{u}(p_1, \lambda_1) i\Gamma_{\mu_1 \nu_1}^{(\mathbb{P}pp)}(p_1, p_a) u(p_a, \lambda_a) \\ &\quad \times i\Delta_{\mu_1 \nu_1, \alpha_1 \beta_1}^{(\mathbb{P})}(s_1, t_1) i\Gamma_{\alpha_1 \beta_1, \alpha_2 \beta_2, \mu}^{(\mathbb{P}P f_1)}(q_1, q_2) i\Delta_{\alpha_2 \beta_2, \mu_2 \nu_2}^{(\mathbb{P})}(s_2, t_2) \\ &\quad \times \bar{u}(p_2, \lambda_2) i\Gamma_{\mu_2 \nu_2}^{(\mathbb{P}pp)}(p_2, p_b) u(p_b, \lambda_b) \end{aligned}$$



with terms of the effective pomeron propagator and the pomeron-proton vertex

$$i\Delta_{\mu\nu, \kappa\lambda}^{(\mathbb{P})}(s, t) = \frac{1}{4s} \left(g_{\mu\kappa} g_{\nu\lambda} + g_{\mu\lambda} g_{\nu\kappa} - \frac{1}{2} g_{\mu\nu} g_{\kappa\lambda} \right) (-is\alpha'_{\mathbb{P}})^{\alpha_{\mathbb{P}}(t)-1}$$

$$i\Gamma_{\mu\nu}^{(\mathbb{P}pp)}(p', p) = -i3\beta_{\mathbb{P}NN} F_1((p' - p)^2) \left\{ \frac{1}{2} [\gamma_\mu(p' + p)_\nu + \gamma_\nu(p' + p)_\mu] - \frac{1}{4} g_{\mu\nu} (\not{p}' + \not{p}) \right\}$$

$$\alpha_{\mathbb{P}}(t) = \alpha_{\mathbb{P}}(0) + \alpha'_{\mathbb{P}} t, \quad \alpha_{\mathbb{P}}(0) = 1.0808, \quad \alpha'_{\mathbb{P}} = 0.25 \text{ GeV}^{-2}$$

$$\beta_{\mathbb{P}NN} = 1.87 \text{ GeV}^{-1}, \quad F_1(t): \text{ Dirac form factor of the proton}$$

Ewerz, Maniatis, Nachtmann, Ann. Phys. 342 (2014) 31


Absorption effects:

$$\mathcal{M}_{pp \rightarrow pp f_1} = \mathcal{M}_{pp \rightarrow pp f_1}^{\text{Born}} + \mathcal{M}_{pp \rightarrow pp f_1}^{\text{pp-rescattering}}$$

$$\mathcal{M}_{pp \rightarrow pp f_1}^{\text{pp-rescattering}}(s, \vec{p}_{1\perp}, \vec{p}_{2\perp}) = \frac{i}{8\pi^2 s} \int d^2 \vec{k}_\perp \mathcal{M}_{pp \rightarrow pp f_1}^{\text{Born}}(s, \vec{p}_{1\perp} - \vec{k}_\perp, \vec{p}_{2\perp} + \vec{k}_\perp) \mathcal{M}_{pp \rightarrow pp}^{\mathbb{P}\text{-exchange}}(s, -\vec{k}_\perp^2)$$

where \vec{k}_\perp is the transverse momentum carried around the loop

IP IP f1 coupling



coupling Lagrangian $\mathcal{L}^{(IPf_1)}$



“bare” vertex function $i\Gamma_{\kappa\lambda,\rho\sigma,\alpha}^{(IPf_1)}(q_1, q_2) |_{\text{bare}}$



CEP reaction

vertex function supplemented by suitable form factor

$$i\tilde{\Gamma}_{\kappa\lambda,\rho\sigma,\alpha}^{(IPf_1)}(q_1, q_2) = i\Gamma_{\kappa\lambda,\rho\sigma,\alpha}^{(IPf_1)}(q_1, q_2) |_{\text{bare}} \tilde{F}_{IPf_1}(q_1^2, q_2^2, k^2)$$

For the on-shell meson we have set $k^2 = m_{f_1}^2$.

$$\tilde{F}^{(IPf_1)}(t_1, t_2, m_{f_1}^2) = F_M(t_1)F_M(t_2), \quad F_M(t) = \frac{1}{1 - t/\Lambda_0^2}, \quad \Lambda_0^2 = 0.5 \text{ GeV}^2$$

or

$$\tilde{F}^{(IPf_1)}(t_1, t_2, m_{f_1}^2) = \exp\left(\frac{t_1 + t_2}{\Lambda_E^2}\right)$$


where the cutoff constant Λ_E should be adjusted to experimental data

We follow two strategies for constructing coupling Lagrangian:

- (1) **Phenomenological approach.** First we consider a fictitious process:
the fusion of two “real spin 2 pomerons” (or tensor glueballs) of mass m giving an f_1 meson of $J^{PC} = 1^{++}$

$$\mathbb{P}(m, \epsilon_1) + \mathbb{P}(m, \epsilon_2) \rightarrow f_1(m_{f_1}, \epsilon)$$

$\epsilon_{1,2}$: polarisation tensors, ϵ : polarisation vector



We work in the rest system of the f_1 meson:

The spin 2 of these “real pomerons” can be combined to a total spin S ($0 \leq S \leq 4$) and this must be combined with the orbital angular momentum ℓ to give $J^{PC} = 1^{++}$ of the f_1 state.

There are exactly two possibilities: $(\ell, S) = (2, 2)$ and $(4, 4)$.

Corresponding couplings are:

$$\mathcal{L}_{\mathbb{P}\mathbb{P}f_1}^{(2,2)} = \frac{g'_{\mathbb{P}\mathbb{P}f_1}}{32 M_0^2} \left(\mathbb{P}_{\kappa\lambda} \overset{\leftrightarrow}{\partial}_\mu \overset{\leftrightarrow}{\partial}_\nu \mathbb{P}_{\rho\sigma} \right) \left(\partial_\alpha U_\beta - \partial_\beta U_\alpha \right) \Gamma^{(8) \kappa\lambda, \rho\sigma, \mu\nu, \alpha\beta}$$

$$\mathcal{L}_{\mathbb{P}\mathbb{P}f_1}^{(4,4)} = \frac{g''_{\mathbb{P}\mathbb{P}f_1}}{24 \times 32 M_0^4} \left(\mathbb{P}_{\kappa\lambda} \overset{\leftrightarrow}{\partial}_{\mu_1} \overset{\leftrightarrow}{\partial}_{\mu_2} \overset{\leftrightarrow}{\partial}_{\mu_3} \overset{\leftrightarrow}{\partial}_{\mu_4} \mathbb{P}_{\rho\sigma} \right) \left(\partial_\alpha U_\beta - \partial_\beta U_\alpha \right) \Gamma^{(10) \kappa\lambda, \rho\sigma, \mu_1\mu_2\mu_3\mu_4, \alpha\beta}$$

Here $M_0 \equiv 1$ GeV, $g'_{\mathbb{P}\mathbb{P}f_1}, g''_{\mathbb{P}\mathbb{P}f_1}$: dimensionless coupling parameters,

$\mathbb{P}_{\kappa\lambda}$ effective pomeron field, U_α f_1 field, $\overset{\leftrightarrow}{\partial}_\mu = \overset{\rightarrow}{\partial}_\mu - \overset{\leftarrow}{\partial}_\mu$ asymmetric derivative,
and $\Gamma^{(8)}, \Gamma^{(10)}$ are known tensor functions.

(2) **Holographic QCD approach** using the Sakai-Sugimoto model.

There, the $IP IP f_1$ coupling can be derived from the bulk **Chern-Simons (CS) term** requiring consistency of supergravity and the gravitational anomaly.

$$\mathcal{L}^{\text{CS}} = \kappa' U_\alpha \varepsilon^{\alpha\beta\gamma\delta} \mathbb{P}^\mu{}_\beta \partial_\delta \mathbb{P}_{\gamma\mu} + \kappa'' U_\alpha \varepsilon^{\alpha\beta\gamma\delta} \left(\partial_\nu \mathbb{P}^\mu{}_\beta \right) \left(\partial_\delta \partial_\mu \mathbb{P}^\nu{}_\gamma - \partial_\delta \partial^\nu \mathbb{P}_{\gamma\mu} \right)$$

κ' : dimensionless, κ'' : dimension GeV^{-2}

Sakai, Sugimoto, Prog. Theor. Phys. 113 (2005) 843; 114 (2005) 1083,
Leutgeb, Rebhan, PRD 101 (2020) 114015

For our fictitious reaction with real pomerons there is strict equivalence $\mathcal{L}^{\text{CS}} \hat{=} \mathcal{L}^{(2,2)} + \mathcal{L}^{(4,4)}$ if the couplings satisfy:

$$g'_{IP IP f_1} = -\kappa' \frac{M_0^2}{k^2} - \kappa'' \frac{M_0^2(k^2 - 2m^2)}{2k^2}$$

$$g''_{IP IP f_1} = \kappa'' \frac{2M_0^4}{k^2}$$

where k^2 is invariant mass squared of the resonance f_1 .





For the CEP reaction

the pomerons have invariant mass squared $t_1, t_2 < 0$ instead of m^2 and, in general, $t_1 \neq t_2$.

Replacing above $2m^2 \rightarrow t_1 + t_2$ we expect for small $|t_1|$ and $|t_2|$ still approximate equivalence to hold.

This is confirmed by explicit numerical studies.

Comparison with experimental results from WA102@CERN



Data: D. Barberis et al. (WA102 Collaboration), PLB 440 (1998) 225

$$f_1(1285) \quad \left| \begin{array}{l} \sqrt{s} = 29.1 \text{ GeV}, |x_{F,M}| \leq 0.2 \\ \sigma_{\text{exp}} = (6919 \pm 886) \text{ nb} \end{array} \right.$$

Phenomenological approach

← (l,S) = (2,2) term only

$$|g'_{\mathbb{P}\mathbb{P}f_1}| = 4.89$$

← (l,S) = (4,4) term only

$$|g''_{\mathbb{P}\mathbb{P}f_1}| = 10.31$$


- We get a reasonable description of WA102 data with $\Lambda_E = 0.7$ GeV
- Absorption effects included
 $\langle S^2 \rangle = \sigma_{\text{abs}} / \sigma_{\text{Born}} \approx 0.5-0.7$
 depending on the kinematics

Comparison with experimental results from WA102@CERN


Comparison with data from: A. Kirk (WA102 Collaboration), Nucl. Phys. A 663 (2000) 608

The theoretical results have been normalized to the mean value of the number of events

Phenomenological approach



- An almost 'flat' distribution at large values of $|t_1 - t_2|$ can be observed
→ absorption effects play a significant role there,
large damping of cross section at higher values of ϕ_{pp}
- It seems that the $(\ell,S) = (4,4)$ term best reproduces the shape of the WA102 data



Holographic QCD approach

← Fit to WA102 data using the Chern-Simons (CS) coupling.


The relation between the (l,S) and CS forms of the couplings:

With $\chi' = -8.88$, $\chi''/\chi' = -1.0 \text{ GeV}^{-2}$

and setting $t_1 = t_2 = -0.1 \text{ GeV}^2$

we get: $g'_{PPf_1} = 0.42$, $g''_{PPf_1} = 10.81$

This CS coupling corresponds practically to a pure $(l,S) = (4,4)$ coupling!



The prediction for χ''/χ' obtained in the Sakai-Sugimoto model:

$$\chi''/\chi' = -5.631/M_{KK}^2 = -(6.25, 3.76, 2.44) \text{ GeV}^{-2}$$


for $M_{KK} = (949, 1224, 1519) \text{ MeV}$

Usually M_{KK} (Kaluza-Klein mass scale) is fixed by matching the mass of the lowest vector meson to that of the physical ρ meson, leading to $M_{KK} = 949 \text{ MeV}$.

However, this choice leads to **tensor glueball mass** which is too low, $M_T \approx 1.5 \text{ GeV}$.

The standard pomeron trajectory corresponds to $M_T \approx 1.9 \text{ GeV}$, whereas lattice gauge theory indicates $M_T \approx 2.4 \text{ GeV}$.

Predictions for the LHC experiments



- The contribution with $\chi''/\chi' = -6.25 \text{ GeV}^{-2}$ gives a significantly different shape
- The absorption effects are included, $\langle S^2 \rangle \approx 0.35$. They decrease the distrib. mostly at higher values of ϕ_{pp} and at smaller values of $p_{t,M}$ (and also $|t|$). This could be tested in ATLAS-ALFA experiment when both protons are measured

Cross sections in μb for $pp \rightarrow pp f_1(1285)$ for $\sqrt{s} = 13 \text{ TeV}$:

Contribution	Parameters $\Lambda_E = 0.7 \text{ GeV}$,	$ y_{f_1} < 1.0$	$ y_{f_1} < 2.5$	$ y_{f_1} < 2.5$, $0.17 < p_{y,p} < 0.50 \text{ GeV}$	$2.0 < y_{f_1} < 4.5$
$(l, S) = (2, 2)$	$g'_{PPf_1} = 4.89$	14.8	37.5	6.46	18.9
$(l, S) = (4, 4)$	$g''_{PPf_1} = 10.31$	13.8	34.0	6.06	18.1
(χ', χ'')	$\chi''/\chi' = -6.25 \text{ GeV}^{-2}$	18.6	45.8	7.14	23.1
(χ', χ'')	$\chi''/\chi' = -2.44 \text{ GeV}^{-2}$	17.5	43.4	7.10	22.1
(χ', χ'')	$\chi''/\chi' = -1.0 \text{ GeV}^{-2}$	16.6	41.0	7.09	20.5

Predictions for the LHC experiments

- One of the most prominent decay modes of the $f_1(1285)$ is $f_1(1285) \rightarrow \pi^+\pi^-\pi^+\pi^-$
- There $f_1(1285)$ and $f_2(1270)$ are close in mass.

We obtain for $\sqrt{s} = 13$ TeV and $|y_M| < 2.5$:

$$\sigma_{pp \rightarrow pp f_1(1285)} \times \mathcal{BR}(f_1(1285) \rightarrow 2\pi^+2\pi^-) = 34.0 \mu\text{b} \times 0.112 = 3.8 \mu\text{b}$$

$$\sigma_{pp \rightarrow pp f_2(1270)} \times \mathcal{BR}(f_2(1270) \rightarrow 2\pi^+2\pi^-) = 11.3 \mu\text{b} \times 0.028 = 0.3 \mu\text{b} \leftarrow \text{CEP of } f_2(1270): \text{Lebiedowicz et al., PRD 93 (2016) 054015, PRD 101 (2020) 034008}$$

As the $f_1(1285)$ has a much narrower width than the $f_2(1270)$ it would be seen in the $M(4\pi)$ distribution as a peak on top of broader $f_2(1270)$ and of the continuum background

- $f_1(1285)$ is [seen in the preliminary ATLAS-ALFA results](#) for $pp \rightarrow pp\pi^+\pi^-\pi^+\pi^-$ at $\sqrt{s} = 13$ TeV and for $|\eta_\pi| < 2.5, p_{t,\pi} > 0.1$ GeV, $\max(p_{t,\pi}) > 0.2$ GeV, 0.17 GeV $< |p_{y,p}| < 0.5$ GeV

[\[R. Sikora, CERN-THESIS-2020-235\]](#)

- Theoretical studies of the reaction $pp \rightarrow pp 4\pi$ including both the resonances and continuum contributions within the tensor-pomeron approach \rightarrow in progress:

4π production via the intermediate $\sigma\sigma$ and $\rho\rho$ states: [Lebiedowicz, Nachtmann, Szczurek, PRD 94 \(2016\) 034017](#)

4π continuum: [Kycia, Lebiedowicz, Szczurek, Turnau, PRD 95 \(2017\) 094020](#)

$f_1(1285)$ production: [Lebiedowicz, Leutgeb, Nachtmann, Rebhan, Szczurek, PRD 102 \(2020\) 114003](#)

using GenEx MC generator for exclusive reactions and DECAY MC library for the decay of a particle with ROOT compatibility:

GenEx MC, [Kycia, Chwastowski, Staszewski, Turnau, Commun. Comput. Phys. 24 \(2018\) 860](#)

DECAY MC, [Kycia, Lebiedowicz, Szczurek, arXiv: 2011.14750 \[hep-ph\], in print Commun. Comput. Phys.](#)

Conclusions

- We have given [predictions for forthcoming experiments with HADES and PANDA at FAIR](#). We have performed feasibility studies and estimated that a 30-days measurement with HADES should allow to identify $f_1(1285)$ meson in the $\pi^+\pi^-\eta$ channel. From such experiments we will learn more on the production mechanism, in particular, about the $\rho\rho f_1$ and $\omega\omega f_1$ coupling strengths.
- We have discussed in detail [CEP of \$f_1\(1285\)\$ meson](#) in pp collisions at high energies in the [tensor-pomeron approach](#). Different forms of the [IP IP \$f_1\$ coupling](#) are possible. Tests of the Sakai-Sugimoto model are possible.
- We obtain a [good description of the WA102 data for the \$pp \rightarrow pp f_1\(1285\)\$ reaction](#) assuming that the reaction is dominated by IP exchange. We have given [predictions for experiments at the LHC](#). We have included - very important - absorptive corrections.

Experimental studies of single meson CEP reactions will give many $IP IP M$ coupling parameters. Their theoretical calculation is a challenging problem of nonperturbative QCD.

Thank you for your attention

Results (HADES and PANDA)

Other decay channels?




PDG: $\mathcal{BR}(f_1(1285) \rightarrow \rho^0 \gamma) = (6.1 \pm 1.0) \%$

CLAS: $\mathcal{BR}(f_1(1285) \rightarrow \rho^0 \gamma) = (2.5^{+0.7}_{-0.8}) \%$

- For the 4π channel it may be difficult to identify the $f_1(1285)$ due to large continuum background e.g. $pp \rightarrow N(1440)N(1440) \rightarrow N\pi\pi N\pi\pi$
 \rightarrow we have found that fusion mechanisms for the $\rho^0\rho^0$ production: π^0 - ω - π^0 and ω - π^0 - ω exchanges (treated with exact $2 \rightarrow 4$ kinematics) give much smaller background cross sections

- The $\rho^0\gamma$ channel should be much better suited. There, however, dominant background channel $pp\pi^+\pi^-\pi^0$ is of the order of 2 mb [1] and ρ^0 is so broad that it will not provide sufficient reductions (as it is the case in η decay channel)

[1] G. Alexander et al., Phys. Rev. 154 (1967) 1284



The $\pi\pi$ -continuum contribution is larger than the VV -continuum term. In both cases the f_1 resonance is clearly visible, even without the reggeization (green lines) in the continuum processes.

We get: for $\sqrt{s} = 3.46 \text{ GeV}$: $\sigma_{pp \rightarrow pp(f_1 \rightarrow \rho^0 \gamma)} = 5.38 \text{ nb}$
 for $\sqrt{s} = 5.0 \text{ GeV}$: $\sigma_{p\bar{p} \rightarrow p\bar{p}(f_1 \rightarrow \rho^0 \gamma)} = 62.86 \text{ nb} \leftarrow 10 \times \text{larger}$

This result makes us rather optimistic that an experimental study of the f_1 in the $\rho\gamma$ decay channel should be possible at PANDA@FAIR.

For our exploratory study we have neglected interference effects between the background $p\gamma$ and the signal $f_1 \rightarrow \rho\gamma$ processes.

We have also neglected the background processes due to bremsstrahlung of γ and ρ^0 from the nucleon lines.